

Is Your Patient Breathing? Ventilation in the OR

Mark J. Haffey MSN, CRNA, APN
TANA Fall Meeting
Oct. 20, 2018



Disclaimer

- ◆ I have no actual or potential conflict of interest in relation to this program/presentation.

THERE'S A LITTLE AUTOPEEP... HE NEEDS
THE FLOW UPPED TO 65... THE ET TUBE CAN
COME BACK 2cm... HIS FOLEY BAG IS FULL...
AND HE WANTS TO WATCH THE LIONS GAME.



TED WAS THE MASTER OF INTERPRETING
VENTILATOR GRAPHICS.

Respiratory Physiology

Ventilation


- ◆ **Inspiration:** Act of drawing air into the lungs, the contraction of muscles creating a reduced pressure in the chest causing lungs to expand and air to flow inward.
- ◆ **Expiration:** Act of breathing out or expelling air out of the lungs. The release of carbon dioxide from the body.
- ◆ **Oxygenation:** The addition of oxygen to any system. Specifically used to describe interventions that provide greater oxygen supply to the lungs and thus the circulation

Ventilatory Drive

- ◆ Chemoreceptors
 - ◆ Carotid Bodies
 - ◆ Stimulated by a ↓ PaO₂, ↓ pH, and hypoperfusion to the receptor.
 - ◆ Primarily Ventilatory Response
 - ◆ Aortic Bodies
 - ◆ Stimulated by a ↓ PaO₂, and hypoperfusion to the receptor
 - ◆ Primarily Circulatory Response
 - ◆ Central Chemoreceptors
 - ◆ Medullary Centers
 - ◆ Located near Cranial Nerve IX and X
 - ◆ Stimulated by ↑ PaCO₂, H⁺, and HCO₃

V/Q: What is the Mismatch?

- ◆ Ratio of Ventilation to Perfusion
- ◆ Physiology that leads to inability for oxygen to be delivered to the alveoli or decrease in perfusion leading to an inability of O₂/CO₂ exchange to take place in the pulmonary system.



Conditions that lead to V/Q mismatch

◆ Decreased V/Q Mismatch

- ◆ Asthma
- ◆ Chronic Bronchitis
- ◆ Pulmonary Edema
- ◆ Airway obstruction

◆ Increased V/Q Mismatch


- ◆ Pulmonary Embolism
- ◆ Emphysema

No Ventilation but adequate perfusion

- a. Pulmonary Embolism
- b. Shunt.

T/F No Perfusion but adequate ventilation = Dead Space

Physiologic V/Q




Patient Positioning: Friend or Foe


- ◆ Trendelenburg:
 - ◆ Cephalad shift of abdominal contents
 - ◆ Decreased ability of the diaphragm to function
 - ◆ Ventilation Zone 3 > Zone 2 > Zone 1
 - ◆ Decreased Compliance
- ◆ Reverse Trendelenburg
 - ◆ Increased Compliance
 - ◆ Caudad shift of abdominal contents
 - ◆ Ventilation Zone 1 > Zone 2 > Zone 3

Ventilator History

Intermittent Negative Pressure Ventilator



Diaphragmatic Ventilator



Ventilator: What is Inside?

- ◆ Bellows: GE (formerly Datex-Omeda)
 - ◆ Driving Gas compresses bellows forcing gas by vaporizers and into the patient.
 - ◆ Driving gas is usually oxygen, sometimes air
- ◆ Piston: Drager Apollo or Fabius GS
 - ◆ Do not require a driving gas
 - ◆ Electric Motor moves a piston thus creating pressure to move air into a patients lungs
- ◆ Turbine: Drager Perseus
 - ◆ Electric motor drives a blower which creates inspiratory pressure and flow
 - ◆ Most efficient ventilator

**Maintain Peak Airway
Pressure <30 cm H₂O**

**Maintain EtCO₂ 32-37.
(exception Neuro cases
where EtCO₂ is desired
<27.)**

**Goals for
Adequate
Ventilation**

**Decrease potential for
Atelectasis**

SaO₂ > 97%


Case Study #1

- ◆ 43 y/o F for Robotic Vaginal Hysterectomy w/BSO secondary to Tumor.
- ◆ PMHx:
 - ◆ BMI 48.2
 - ◆ Obstructive Sleep Apnea (CPAP at home)
 - ◆ HTN
 - ◆ NIDDM
- ◆ Position: Steep Trendelenburg
- ◆ Anesthesia: General Anesthesia
 - ◆ Paralyzed
 - ◆ ETT
- ◆ Ventilation: WHAT WOULD YOU DO?

Mechanical Ventilation in the OR

- ◆ **Positive Pressure Ventilator**
- ◆ **Used to adequately ventilate Patients**
 - ◆ Decreased lung compliance
 - ◆ Increased Airway Resistance
 - ◆ Absence of ventilation drive secondary to physiologic changes
- ◆ **Multiple ventilation modes**
- ◆ **Computer-Driven with active patient monitoring and feedback systems to enable adequate ventilation.**

Why Do We Need Mechanical Ventilation?



Reasons for Mechanical Ventilation

◆ **Surgical Procedure**

- ◆ Intrathoracic or Intra-abdominal
- ◆ ENT
- ◆ Muscle Relaxation required for adequate exposure or decreased risk of complication.

◆ **Inability to adequately oxygenate a patient.**

- ◆ Patient physiology
- ◆ Pulmonary Disease
- ◆ Patient position

Anesthesia Machine Ventilator Modes

Volume Control

- Selected Specific Volume
- Rate Selected
- Independent of patient effort or response

Pressure Controlled

- Selected Pressure
- Variable Volume
- Rate Selected

SIMV

- Spontaneous Breaths allowed
- Preset Rate
- Preset Pressure

Anesthesia Ventilator Modes (cont.)

Assist Control Ventilation

- Intermittent positive pressure
- Pt initiates ventilator to deliver a pre-determined tidal volume.
- Back-up mode should patient be unable to initiate a breath

PSV: Pressure Support Ventilation

- Preset level of positive airway pressure
- Patient determines ventilatory rate
- Designed to increase the volume of the patients spontaneous breath
- Decreases overall work of breathing especially diaphragmatic fatigue

Extended Mandatory Minute Ventilation

- Preset Minute volume of gas either from a positive pressure breath or a spontaneous breath
- Preset mechanical rate and volume
- The goal is have patients use more of their respiratory muscles

Adjunct Ventilator Settings

◆ I:E

- ◆ Ratio of Inspiration to Expiration
 - ◆ Time Inspired (PCV)
 - ◆ ↓PAP when using I:E to increase inspiratory time
 - ◆ ↑ expiratory time to allow adequate CO₂ removal in COPD patient
 - ◆ Inverse Ratio Ventilation

◆ PEEP

- ◆ Positive Pressure designed to keep alveoli open throughout the respiratory cycle.
- ◆ Intrinsic vs. Extrinsic

Hyperoxemia: Is it important

- ◆ Suzuki, S. et al. Anesthesiology 2018: Randomized Study of 1786 patients
 - ◆ Results showed that in 92% of patients O₂ was maintained between .32 and .6. 1% of patients had O₂<.3 and 7% of patients had O₂> .7
 - ◆ Of these patients 83% had hyperoxemia with 32% of patients having significant hyperoxemia
- ◆ Applegate, et al.: Intraoperative Hyperoxemia: An Unnecessary Evil
 - ◆ Mean PaO₂ was 206mmHg with values up to 534mmHg of O₂
 - ◆ Excessive O₂ can lead to increase in reactive oxidative agents leading to cell damage, and dysfunction
 - ◆ Recommended to have O₂ below 150mmHg

Pressure Controlled Ventilation

- ◆ Volume is determined by preset pressure
- ◆ Time Triggered ventilatory cycle
- ◆ Ability to be delivered in IMV or ACV modes
- ◆ Has a Decelerating Flow Pattern which improves distribution of gases
- ◆ Ideal for patient with decreased compliance due to
 - ◆ Body habitus
 - ◆ Pt position
 - ◆ Surgical Case

Must set Maximum Tidal Volume, especially laparoscopic cases!

VENTILATOR GEM

- ◆ **Pressure Control with Volume Guarantee (PCV-VG)**
 - ◆ Volume and Pressure are preset
 - ◆ Preset Volume with a Max PAP!
 - ◆ Ventilator adjusts P_{insp} to achieve preset TV breath by breath
 - ◆ Key Advantages:
 - ◆ Controlled Peak Inspiratory Pressure
 - ◆ Control of Minute Ventilation
 - ◆ Preset Volume delivered each breath at the minimum necessary pressure

SIMV


- ◆ Synchronized Intermittent Mandatory Ventilation:
 - ◆ Combination of ACV and Spontaneous Ventilation
- ◆ Baseline Respiratory Rate that allows patient to breath at their own rate between ventilator initiated breaths
- ◆ Preset Inspiratory pressure for intermittent Mandatory breaths
- ◆ Ventilator attempts to synchronize mandatory breaths with patient initiated breaths.
- ◆ **PSV** is used in conjunction with SIMV
 - ◆ Pt sets own respiratory rate with minimum pressure applied for each breath

Assist Controlled Ventilation

- ◆ Intermittent positive-pressure ventilation mode in which the patient creates a sub-baseline pressure in the inspiratory limb which then triggers the ventilator to deliver a predetermined tidal volume.
 - ◆ Back-up control mode should respiratory rate drop below a preset level.
- ◆ Every breath is same volume whether patient initiates or the ventilator delivers
- ◆ Not for patients with a rapid respiratory rate

What does the Research Say

- ◆ Gajic, et. al: Patients ventilated with TV >12% IBW were 25% more likely to have ALI or ARDS
- ◆ Choi et al.: Study noted that patients that were ventilated with 12 ml/kg at 5 hours showed significant procoagulant changes indicative of Acute Lung Injury when compared to Lung Volumes of 6ml/kg with 10cm PEEP.
- ◆ Research done by the International Anesthesia Research Society shows PEEP increases intrathoracic pressure and the effects are amplified in patients suffering hypovolemia or cardiac dysfunction.. Does not recommend that SaO₂ be used as an endpoint for PEEP



Pneumonia

Pneumothorax

Acute Lung Injury

Atelectasis

Complication
of Mechanical
Ventilation

ARDS

Keys to overcoming Ventilatory Complications

💧 **Pre-oxygenation**

- 💧 ↓ Hypoxia
- 💧 ↑ FRC
- 💧 ↓ Reabsorption Atelectasis

💧 **Obesity**

- 💧 ↓ FRC
- 💧 37% of closed claims related to difficult intubation involved obese patients.

💧 **Intermittent Alveolar Recruitment Breaths**

- 💧 **Essential** that pre-oxygenation with a Facemask is practiced.

Pre-Existing Conditions that Compromise Ventilation

◆ COPD

- ◆ Emphysema
- ◆ Asthma
- ◆ Chronic Bronchitis

◆ OSA

◆ Tracheal Stenosis

◆ Previous Lung Resection

◆ ARDS

The “Not-So” wanted consequences

- ◆ Hypotension
 - ◆ Increased Intrathoracic Pressure causing Decreased Venous Return
 - ◆ Positive Pressure Ventilation
 - ◆ Potentiated w/Insufflation for Laparoscopic Procedures
- ◆ ↓Cardiac Output
- ◆ Airway
 - ◆ Tracheal Scarring as a result of intubation
 - ◆ Negative Pressure Pulmonary Edema
 - ◆ Vocal Cord Damage

Case Study #2

- ◆ 38 y/o M for Laparoscopic Robotic Gastric Bypass
- ◆ BMI 58.2
- ◆ PMHx:
 - ◆ HTN
 - ◆ NIDDM
 - ◆ OSA
 - ◆ Gout
- ◆ PSHx:
 - ◆ Appy
 - ◆ Cholecystectomy
 - ◆ Shoulder Arthroscopy
- ◆ What is your Anesthesia and Ventilation Plan?

Case Study #2 (cont.)

- ◆ Position: Reverse Trendelenburg
- ◆ Endoscopy performed(? Surgeon Dependent)
- ◆ Induction:
 - ◆ Lidocaine
 - ◆ Propofol
 - ◆ Succinylcholine
 - ◆ Remi-Fentanyl
 - ◆ (.75mcg/kg/min followed with a maintenance of .2mcg/kg/min)
 - ◆ Dilaudid
 - ◆ 1mg at the end of the procedure
- ◆ ERAS Protocol if surgeons are on board.
- ◆ Minimal Post-op Pain secondary to local anesthetic

How are you going to ventilate this patient?

Optimizing Ventilation

- ◆ Tidal Volume (6 – 8ml/kg) although lower volumes are ideal
- ◆ PEEP 5cm H₂O
- ◆ Respiratory Rate
 - ◆ Maintain pCO₂ 32-37 (homeostatic state)
- ◆ Peak Airway Pressure < 30cm H₂O
 - ◆ ARDS
 - ◆ Utilize I:E Ratio
- ◆ Intermittent Recruitment Breaths
 - ◆ Manually Insufflate to a PAP 40-50.
 - ◆ Every 30min.

Must be able to adapt for DECREASED Compliance and INCREASED Resistance

PSV PRO “Unique to Anesthesia”

- ◆ Optimization of the spontaneously breathing patient.
- ◆ Respiratory Rate driven by patient
 - ◆ Should patient not breath for a preset time (30sec) SIMV mode ventilation is initiated.
 - ◆ Protective mode with preset rate and pressure for SIMV mode
- ◆ Pressure supported breath to achieve adequate TV for each patient initiated breath.
- ◆ Research shows it is a superior mode to SIMV when weaning a patient from mechanical ventilation.

Summary

- ◆ Mechanical Ventilation is a standard of care in General Anesthesia
- ◆ Selection of Ventilation mode should be determined by
 - ◆ Surgical Case
 - ◆ Patient position
 - ◆ Pre-existing respiratory disease and function
 - ◆ Patient BMI
- ◆ Prevent atelectasis by using low TV and low PEEP.
- ◆ Do not be afraid to use different ventilator modes and I:E Ratio to maximize ventilation.

Questions??



References

- Barash, PG et al. (1997) *Clinical Anesthesia-3rd Edition*. Philadelphia, Lippincott-Raven Publishers
- Bristle, TJ et. al “Anesthesia and Critical Care Ventilator Modes: Past, Present, and Future”, AANA Journal. Oct 2014 82(5) p387-400.
- Futier, E. et.al “Mechanical Ventilation in Abdominal Surgery” Annales Francaoise d’ Aneshtesie et de reanimation. 2014 33(7-8) p. 472-74.
- Habre, W. “Complex Ventilation Stratigies in Paediatric Anesthesia” European Society of Anesthesiology. 2010 pg 1-5.
- International Anesthesia Research Society, (2016) Modes of Mechanical Ventilation.

References (Cont.)

- ◆ Seok, YS et.al “Volume controlled versus Pressure Controlled Ventilation-Volume Guaranteed Mode during One-Lung Ventilation”, Korean Journal of Anesthesiology. Oct 2014 67(4) p258-63.
- ◆ Shubhangi, A et. Al “Ventilatory Strategies in Trauma Patients”, Journal of Emergencies, Trauma, and Shock. Jan-Mar 2014 7(1) p25-31.
- ◆ Slinger, P. “Don’t Make Things Worse with your Ventilator Settings: How You Manage the Lungs During Perioperative Period Affects Postoperative Outcomes” International Anesthesia Research Society Review Course. 2013 pg 38-46.